Stirling P, Tahir M, Atkinson HD. The Limitations of Gram-stain Microscopy of Synovial Fluid in Concomitant Septic and Crystal Arthritis. Curr Rheumatol Rev 2018; 14:255. Boumans D, Hettema ME, Vonkeman HE, et al. The added value of synovial fluid centrifugation for monosodium urate and calcium pyrophosphate crystal detection. Clin Rheumatol 2017; 36:1599. Meenagh GK, Patton J, Kynes C, Wright GD. A randomised controlled trial of intraarticular corticosteroid injection of the carpometacarpal joint of the thumb in osteoarthritis. Ann Rheum Dis 2004; 63:1260. Stahl S, Karsh-Zafrir I, Ratzon N, Rosenberg N. Comparison of intraarticular injection of depot corticosteroid and hyaluronic acid for treatment of degenerative trapeziometacarpal joints. J Clin Rheumatol 2005; 11:299. Geirsson AJ, Statkevicius S, VÃkingsson A. Septic arthritis in Iceland 1990-2002: increasing incidence due to iatrogenic infections. Ann Rheum Dis 2008; 67:638. Sibbitt WL Jr, Peisajovich A, Michael AA, et al. Does sonographic needle guidance affect the clinical outcome of intraarticular injections? J Rheumatol 2009; 36:1892. Cunnington J, Marshall N, Hide G, et al. A randomized, double-blind, controlled study of ultrasound-guided corticosteroid injection into the joint of patients with inflammatory arthritis. Arthritis Rheum 2010; 62:1862. Charalambous CP, Tryfonidis M, Sadiq S, et al. Septic arthritis following intra-articular steroid injection of the knee--a survey of current practice regarding antiseptic technique used during intra-articular steroid injection of the knee. Clin Rheumatol 2003; 22:386. Mimoz O, Karim A, Mercat A, et al. Chlorhexidine compared with povidone-iodine as skin preparation before blood culture. A randomized, controlled trial. Ann Intern Med 1999; 131:834. Polishchuk D, Gehrmann R, Tan V. Skin sterility after application of ethyl chloride spray. J Bone Joint Surg Am 2012; 94:118. Kirschke DL, Jones TF, Stratton CW, et al. Outbreak of joint and soft-tissue infections associated with injections from a multiple-dose medication vial. Clin Infect Dis 2003; 36:1369. Mattner F, Gastmeier P. Bacterial contamination of multiple-dose vials: a prevalence study. Am J Infect Control 2004; 32:12. Motamedifar M, Askarian M. The prevalence of multidose vial contamination by aerobic bacteria in a major teaching hospital, Shiraz, Iran, 2006. Am J Infect Control 2009; 37:773. Schaefer MK, Shehab N, Perz JF. Calling it 'multidose' doesn't make it so: inappropriate sharing and contamination of parenteral medication vials. Am J Infect Control 2010; 38:580. Weitoft T, Uddenfeldt P. Importance of synovial fluid aspiration when injecting intra-articular corticosteroids. Ann Rheum Dis 2000; 59:233. Thumboo J, O'Duffy JD. A prospective study of the safety of joint and soft tissue aspirations and injections in patients taking warfarin sodium. Arthritis Rheum 1998; 41:736. Salvati G, Punzi L, Pianon M, et al. [Frequency of the bleeding risk in patients receiving warfarin submitted to arthrocentesis of the knee]. Reumatismo 2003; 55:159. Ahmed I, Gertner E. Safety of arthrocentesis and joint injection in patients receiving anticoagulation at therapeutic levels. Am J Med 2012; 125:265. Yui JC, Preskill C, Greenlund LS. Arthrocentesis and Joint Injection in Patients Receiving Direct Oral Anticoagulants. Mayo Clin Proc 2017; 92:1223. Bashir MA, Ray R, Sarda P, et al. Determination of a safe INR for joint injections in patients taking warfarin. Ann R Coll Surg Engl 2015; 97:589. Conway R, O'Shea FD, Cunnane G, Doran MF. Safety of joint and soft tissue injections in patients on warfarin anticoagulation. Clin Rheumatol 2013; 32:1811. Wise, CM. Aspiration and injection of joints in soft tissue. In: Textbook of Rheumatology, 8th Edition, Firestein, G, Budd, R, Harris, T, et al. (Eds), WB Saunders, Philadelphia 2009. p.721. Strobel G, Schwarz R, Heppt P, et al. [Use of an improved transmission ultrasound camera for diagnosis of rheumatic joint diseases]. Z Rheumatol 1993;

## 52:114. Hirsch G, O'Neill T, Kitas G, Klocke R. Distribution of effusion in knee arthritis as measured by high-resolution ultrasound. Clin Rheumatol 2012; 31:1243.

Grant's Atlas of Anatomy, 14th ed, Agur A, Dalley A (Eds), Lippincott Williams & Wilkins, Philadelphia 2017. Thomson CJ, Lalonde DH. Randomized double-blind comparison of duration of anesthesia among three commonly used agents in digital nerve block. Plast Reconstr Surg 2006; 118:429. Alhelail M, Al-Salamah M, Al-Mulhim M, Al-Hamid S. Comparison of bupivacaine and lidocaine with epinephrine for digital nerve blocks. Emerg Med J 2009; 26:347. Valvano MN, Leffler S. Comparison of bupivacaine and lidocaine/bupivacaine for local anesthesia/digital nerve block. Ann Emerg Med 1996; 27:490. Altinyazar HC, Ozdemir H, Koca R, et al. Epinephrine in digital block: color Doppler flow imaging. Dermatol Surg 2004; 30:508. Prabhakar H, Rath S, Kalaivani M, Bhanderi N. Adrenaline with lidocaine for digital nerve blocks. Cochrane Database Syst Rev 2015; :CD010645. Denkler K. A comprehensive review of epinephrine in the finger: to do or not to do. Plast Reconstr Surg 2001; 108:114. Krunic AL, Wang LC, Soltani K, et al. Digital anesthesia with epinephrine: an old myth revisited. J Am Acad Dermatol 2004; 51:755. Muck AE, Bebarta VS, Borys DJ, Morgan DL. Six years of epinephrine digital injections: absence of significant local or systemic effects. Ann Emerg Med 2010; 56:270. Chowdhry S, Seidenstricker L, Cooney DS, et al. Do not use epinephrine in digital blocks: myth or truth? Part II. A retrospective review of 1111 cases. Plast Reconstr Surg 2010; 126:2031. Mohan PP. Towards evidence based emergency medicine: best BETs from the Manchester Royal Infirmary. Epinephrine in digital nerve block. Emerg Med J 2007; 24:789. Lalonde D, Bell M, Benoit P, et al. A multicenter prospective study of 3,110 consecutive cases of elective epinephrine use in the fingers and hand: the Dalhousie Project clinical phase. J Hand Surg Am 2005; 30:1061. Wilhelmi BJ, Blackwell SJ, Miller J, et al. Epinephrine in digital blocks: revisited. Ann Plast Surg 1998; 41:410. Thomson CJ, Lalonde DH, Denkler KA, Feicht AJ. A critical look at the evidence for and against elective epinephrine use in the finger. Plast Reconstr Surg 2007; 119:260. Sonohata M, Nagamine S, Maeda K, et al. Subcutaneous single injection digital block with epinephrine. Anesthesiol Res Pract 2012; 2012:487650. Fitzcharles-Bowe C, Denkler K, Lalonde D. Finger injection with high-dose (1:1,000) epinephrine: Does it cause finger necrosis and should it be treated? Hand (N Y) 2007; 2:5. Ilicki J. Safety of Epinephrine in Digital Nerve Blocks: A Literature Review. J Emerg Med 2015; 49:799. Zilinsky I, Bar-Meir E, Zaslansky R, et al. Ten commandments for minimal pain during administration of local anesthetics. J Drugs Dermatol 2005; 4:212. Bartfield JM, Ford DT, Homer PJ. Buffered versus plain lidocaine for digital nerve blocks. Ann Emerg Med 1993; 22:216. Cornelius P, Kendall J, Meek S, Rajan R. Alkalinisation of lignocaine to reduce the pain of digital nerve blockade. J Accid Emerg Med 1996; 13:339. Waldbillig DK, Quinn JV, Stiell IG, Wells GA. Randomized double-blind controlled trial comparing room-temperature and heated lidocaine for digital nerve block. Ann Emerg Med 1995; 26:677. Browne J, Fung M, Donnelly M, Cooney C. The use of EMLA reduces the pain associated with digital ring block for ingrowing toenail correction. Eur J Anaesthesiol 2000; 17:182. Serour F, Ben-Yehuda Y, Boaz M. EMLA cream prior to digital nerve block for ingrown nail surgery does not reduce pain at injection of anesthetic solution. Acta Anaesthesiol Scand 2002; 46:203. Morrison WG. Transthecal digital block. Arch Emerg Med 1993; 10:35. Hill RG Jr, Patterson JW, Parker JC, et al. Comparison of transthecal digital block and traditional digital block for anesthesia of the finger. Ann Emerg Med 1995; 25:604. Low CK, Vartany A, Engstrom

JW, et al. Comparison of transthecal and subcutaneous single-injection digital block techniques. J Hand Surg Am 1997; 22:901. Keramidas EG, Rodopoulou SG, Tsoutsos D, et al. Comparison of transthecal digital block and traditional digital block for anesthesia of the finger. Plast Reconstr Surg 2004; 114:1131. Sonohata M, Asami A, Ogawa K, et al. Single injection digital block: is a transthecal injection necessary? J Hand Surg Eur Vol 2009; 34:94. Antevy PM, Zuckerbraun NS, Saladino RA, Pitetti RD. Evaluation of a transthecal digital nerve block in the injured pediatric patient. Pediatr Emerg Care 2010; 26:177. Cannon B, Chan L, Rowlinson JS, et al. Digital anaesthesia: one injection or two? Emerg Med J 2010; 27:533. Hung VS, Bodavula VK, Dubin NH. Digital anaesthesia: comparison of the efficacy and pain associated with three digital nerve block techniques. J Hand Surg Br 2005; 30:581. Brutus JP, Baeten Y, Chahidi N, et al. Single injection digital block: comparison between three techniques. Chir Main 2002; 21:182. Martin SP, Chu KH, Mahmoud I, et al. Double-dorsal versus single-volar digital subcutaneous anaesthetic injection for finger injuries in the emergency department: A randomised controlled trial. Emerg Med Australas 2016; 28:193. Okur OM, Åžener A, Kavakli HÅž, et al. Two injection digital block versus single subcutaneous palmar injection block for finger lacerations. Eur J Trauma Emerg Surg 2017; 43:863. Whiteley B, Rees S. A randomized controlled trial to compare two techniques for partial digital local anesthetic blocks. J Foot Ankle Surg 2010; 49:143. Salasche SJ. Surgery. In: Nails: Diagnosis, Therapy, Surgery, 3rd, Scher RK, Daniel CR (Eds), Saunders, Philadelphia 2005. p.267.

Cohen SP. Epidemiology, diagnosis, and treatment of neck pain. Mayo Clin Proc 2015; 90:284. Grubb SA, Kelly CK. Cervical discography: clinical implications from 12 years of experience. Spine (Phila Pa 1976) 2000; 25:1382. Slipman CW, Plastaras C, Patel R, et al. Provocative cervical discography symptom mapping. Spine J 2005; 5:381. Boden SD, McCowin PR, Davis DO, et al. Abnormal magnetic-resonance scans of the cervical spine in asymptomatic subjects. A prospective investigation. J Bone Joint Surg Am 1990; 72:1178. Teresi LM, Lufkin RB, Reicher MA, et al. Asymptomatic degenerative disk disease and spondylosis of the cervical spine: MR imaging. Radiology 1987; 164:83. Schellhas KP, Smith MD, Gundry CR, Pollei SR. Cervical discogenic pain. Prospective correlation of magnetic resonance imaging and discography in asymptomatic subjects and pain sufferers. Spine (Phila Pa 1976) 1996; 21:300. Bogduk N, Marsland A. On the concept of third occipital headache. J Neurol Neurosurg Psychiatry 1986; 49:775. Lord SM, Barnsley L, Wallis BJ, Bogduk N. Chronic cervical zygapophysial joint pain after whiplash. A placebo-controlled prevalence study. Spine (Phila Pa 1976) 1996; 21:1737. Manchikanti L, Singh V, Rivera J, Pampati V. Prevalence of cervical facet joint pain in chronic neck pain. Pain Physician 2002; 5:243. Gennis P, Miller L, Gallagher EJ, et al. The effect of soft cervical collars on persistent neck pain in patients with whiplash injury. Acad Emerg Med 1996; 3:568. Pain in the neck and arm: a multicentre trial of the effects of physiotherapy, arranged by the British Association of Physical Medicine. Br Med J 1966; 1:253. Aker PD, Gross AR, Goldsmith CH, Peloso P. Conservative management of mechanical neck pain: systematic overview and meta-analysis. BMJ 1996; 313:1291. Devo RA. Drug therapy for back pain. Which drugs help which patients? Spine (Phila Pa 1976) 1996; 21:2840. Peloso P, Gross A, Haines T, et al. Medicinal and injection therapies for mechanical neck disorders. Cochrane Database Syst Rev 2007; :CD000319. Borenstein DG, Korn S. Efficacy of a low-dose regimen of cyclobenzaprine hydrochloride in acute skeletal muscle spasm: results of two

placebo-controlled trials. Clin Ther 2003; 25:1056. Turturro MA, Frater CR, D'Amico FJ. Cyclobenzaprine with ibuprofen versus ibuprofen alone in acute myofascial strain: a randomized, double-blind clinical trial. Ann Emerg Med 2003; 41:818. Bronfort G, Evans R, Anderson AV, et al. Spinal manipulation, medication, or home exercise with advice for acute and subacute neck pain: a randomized trial. Ann Intern Med 2012; 156:1. Mealy K, Brennan H, Fenelon GC. Early mobilization of acute whiplash injuries. Br Med J (Clin Res Ed) 1986; 292:656. Rosenfeld M, Gunnarsson R, Borenstein P. Early intervention in whiplash-associated disorders: a comparison of two treatment protocols. Spine (Phila Pa 1976) 2000; 25:1782. Verhagen AP, Peeters GG, de Bie RA, Oostendorp RA. Conservative treatment for whiplash. Cochrane Database Syst Rev 2001; :CD003338. Chow RT, Johnson MI, Lopes-Martins RA, Bjordal JM. Efficacy of low-level laser therapy in the management of neck pain: a systematic review and meta-analysis of randomised placebo or active-treatment controlled trials. Lancet 2009; 374:1897. Schellingerhout JM, Heymans MW, Verhagen AP, et al. Prognosis of patients with nonspecific neck pain: development and external validation of a prediction rule for persistence of complaints. Spine (Phila Pa 1976) 2010; 35:E827. Shahidi B, Curran-Everett D, Maluf KS. Psychosocial, Physical, and Neurophysiological Risk Factors for Chronic Neck Pain: A Prospective Inception Cohort Study. J Pain 2015; 16:1288. Valenza MC, Valenza G, GonzÃjlez-JimÃ@nez E, et al. Alteration in sleep quality in patients with mechanical insidious neck pain and whiplash-associated neck pain. Am J Phys Med Rehabil 2012; 91:584. Richter M, Ferrari R, Otte D, et al. Correlation of clinical findings, collision parameters, and psychological factors in the outcome of whiplash associated disorders. J Neurol Neurosurg Psychiatry 2004; 75:758. CÃ'té P, Hogg-Johnson S, Cassidy JD, et al. Initial patterns of clinical care and recovery from whiplash injuries: a population-based cohort study. Arch Intern Med 2005; 165:2257. Walton DM, Macdermid JC, Giorgianni AA, et al. Risk factors for persistent problems following acute whiplash injury: update of a systematic review and meta-analysis. J Orthop Sports Phys Ther 2013; 43:31. Gross A, Kay TM, Paquin JP, et al. Exercises for mechanical neck disorders. Cochrane Database Syst Rev 2015; 1:CD004250.

Kucik CJ, Clenney T. Management of epistaxis. Am Fam Physician 2005; 71:305. Petruson B, Rudin R. The frequency of epistaxis in a male population sample. Rhinology 1975; 13:129. Villwock JA, Jones K. Recent trends in epistaxis management in the United States: 2008-2010. JAMA Otolaryngol Head Neck Surg 2013; 139:1279. Kotecha B, Fowler S, Harkness P, et al. Management of epistaxis: a national survey. Ann R Coll Surg Engl 1996; 78:444. Pallin DJ, Chng YM, McKay MP, et al. Epidemiology of epistaxis in US emergency departments, 1992 to 2001. Ann Emerg Med 2005; 46:77. Tomkinson A, Roblin DG, Flanagan P, et al. Patterns of hospital attendance with epistaxis. Rhinology 1997; 35:129. Daniell HW. Estrogen prevention of recurrent epistaxis. Arch Otolaryngol Head Neck Surg 1995; 121:354. Fishpool SJ, Tomkinson A. Patterns of hospital admission with epistaxis for 26,725 patients over an 18-year period in Wales, UK. Ann R Coll Surg Engl 2012; 94:559. Nunez DA, McClymont LG, Evans RA. Epistaxis: a study of the relationship with weather. Clin Otolaryngol Allied Sci 1990; 15:49. Okafor BC. Epistaxis: a clinical study of 540 cases. Ear Nose Throat J 1984; 63:153. Juselius H. Epistaxis. A clinical study of 1,724 patients. J Laryngol Otol 1974; 88:317. Lund VJ, Preziosi P, Hercberg S, et al. Yearly incidence of rhinitis, nasal bleeding, and other nasal symptoms in mature women. Rhinology 2006; 44:26. Manfredini R, Gallerani M, Portaluppi F. Seasonal variation in the

occurrence of epistaxis. Am J Med 2000; 108:759. Pollice PA, Yoder MG. Epistaxis: a retrospective review of hospitalized patients. Otolaryngol Head Neck Surg 1997; 117:49. Schlosser RJ. Clinical practice. Epistaxis. N Engl J Med 2009; 360:784. Alvi A, Joyner-Triplett N. Acute epistaxis. How to spot the source and stop the flow. Postgrad Med 1996; 99:83. Nitu IC, Perry DJ, Lee CA. Clinical experience with the use of clotting factor concentrates in oral anticoagulation reversal. Clin Lab Haematol 1998; 20:363. Lavy J. Epistaxis in anticoagulated patients: educating an at-risk population. Br J Haematol 1996; 95:195. Shah RK, Dhingra JK, Shapshay SM. Hereditary hemorrhagic telangiectasia: a review of 76 cases. Laryngoscope 2002; 112:767. Fuchizaki U, Miyamori H, Kitagawa S, et al. Hereditary haemorrhagic telangiectasia (Rendu-Osler-Weber disease). Lancet 2003; 362:1490. Fiorella ML, Ross D. Henderson KJ. White RI Jr. Outcome of septal dermoplasty in patients with hereditary hemorrhagic telangiectasia. Laryngoscope 2005; 115:301. Liu JK, Gottfried ON, Amini A, Couldwell WT. Aneurysms of the petrous internal carotid artery: anatomy, origins, and treatment. Neurosurg Focus 2004; 17:E13. Chen D, Concus AP, Halbach VV, Cheung SW. Epistaxis originating from traumatic pseudoaneurysm of the internal carotid artery: diagnosis and endovascular therapy. Laryngoscope 1998; 108:326. Hart RG, Pearce LA. In vivo antithrombotic effect of aspirin: dose versus nongastrointestinal bleeding. Stroke 1993; 24:138. Beran M, Petruson B. Occurrence of epistaxis in habitual nose-bleeders and analysis of some etiological factors. ORL J Otorhinolaryngol Relat Spec 1986; 48:297.