Johansen K, Daines M, Howey T, et al. Objective criteria accurately predict amputation following lower extremity trauma. J Trauma 1990; 30:568. de Mestral C, Sharma S, Haas B, et al. A contemporary analysis of the management of the mangled lower extremity. J Trauma Acute Care Surg 2013; 74:597. Johnson BA, Carmack D, Neary M, et al. Operation Iragi Freedom: the Landstuhl Regional Medical Center experience. J Foot Ankle Surg 2005; 44:177. Mackenzie EJ, Fowler CJ. Epidemiology. In: Trauma, 6th ed., Feliciano DV, Mattox KL, Moore EE (Eds), McGraw-Hill Medical, New York 2008. p.25. Nance ML. National Trauma Data Bank Annual Report. 2012. http://www.facs.org/trauma/ntdb/pdf/ntdb-annual-report-2012.pdf (Accessed on October 22, 2013). Finkelstein EA, Corso PS, Miller TR, et al. The incidence and economic burden of injuries in the United States, Oxford University Press, New York 2006. Perkins ZB, Yet B, Glasgow S, et al. Meta-analysis of prognostic factors for amputation following surgical repair of lower extremity vascular trauma. Br J Surg 2015; 102:436. Owens BD, Kragh JF Jr, Wenke JC, et al. Combat wounds in operation Iragi Freedom and operation Enduring Freedom. J Trauma 2008; 64:295. Belmont PJ Jr, Goodman GP, Zacchilli M, et al. Incidence and epidemiology of combat injuries sustained during "the surge" portion of operation Iraqi Freedom by a U.S. Army brigade combat team. J Trauma 2010; 68:204. Fox CJ, Gillespie DL, O'Donnell SD, et al. Contemporary management of wartime vascular trauma. J Vasc Surg 2005; 41:638. Dillingham TR, Pezzin LE, MacKenzie EJ. Incidence, acute care length of stay, and discharge to rehabilitation of traumatic amputee patients: an epidemiologic study. Arch Phys Med Rehabil 1998; 79:279. Meling T, Harboe K, SÃ, reide K. Incidence of traumatic long-bone fractures requiring inhospital management: a prospective age- and gender-specific analysis of 4890 fractures. Injury 2009; 40:1212. Court-Brown CM, Rimmer S, Prakash U, McQueen MM. The epidemiology of open long bone fractures. Injury 1998; 29:529. Court-Brown CM, McBirnie J. The epidemiology of tibial fractures. J Bone Joint Surg Br 1995; 77:417. Schlickewei W, Kuner EH, Mullaji AB, Götze B. Upper and lower limb fractures with concomitant arterial injury. J Bone Joint Surg Br 1992; 74:181. Frykberg ER, Dennis JW, Bishop K, et al. The reliability of physical examination in the evaluation of penetrating extremity trauma for vascular injury: results at one year. J Trauma 1991; 31:502. Franz RW, Shah KJ, Halaharvi D, et al. A 5-year review of management of lower extremity arterial injuries at an urban level I trauma center. J Vasc Surg 2011; 53:1604. White JM, Stannard A, Burkhardt GE, et al. The epidemiology of vascular injury in the wars in Iraq and Afghanistan. Ann Surg 2011; 253:1184. Patel JA, White JM, White PW, et al. A contemporary, 7year analysis of vascular injury from the war in Afghanistan. J Vasc Surg 2018; 68:1872. https://www.facs.org/quality-programs/trauma/atls (Accessed on February 19, 2019). American College of Surgeons Committee on Trauma. Advanced Trauma Life Support (ATLS) Student Course Manual, 9th ed, American College of Surgeons, Chicago 2012. Bulger EM, Snyder D, Schoelles K, et al. An evidence-based prehospital guideline for external hemorrhage control: American College of Surgeons Committee on Trauma. Prehosp Emerg Care 2014; 18:163. Dorlac WC, DeBakey ME, Holcomb JB, et al. Mortality from isolated civilian penetrating extremity injury. J Trauma 2005; 59:217. Swan KG Jr, Wright DS, Barbagiovanni SS, et al. Tourniquets revisited. J Trauma 2009; 66:672. Arrillaga A, Bynoe R, Frykberg ER, Nagy K. EAST Practice Management Guidelines for Penetrating Trauma to the Lower Extremity (2002). http://www.east.org/content/documents/lower extremity isolated arterial injuries from pe netrating trauma.pdf (Accessed on October 22, 2013). Fox N, Rajani RR, Bokhari F, et al. Evaluation and management of penetrating lower extremity arterial trauma: an Eastern

Association for the Surgery of Trauma practice management guideline. J Trauma Acute Care Surg 2012; 73:S315. Committee for Tactical Emergency Medical Care. Tactical Emergency Casualty Care Guidelines: Direct Threat Care. http://c-tecc.org/tactical-emergency-casualtycare-guidelines (Accessed on October 22, 2013). Beekley AC, Sebesta JA, Blackbourne LH, et al. Prehospital tourniquet use in Operation Iraqi Freedom: effect on hemorrhage control and outcomes. J Trauma 2008; 64:S28. Kragh JF Jr, Walters TJ, Baer DG, et al. Survival with emergency tourniquet use to stop bleeding in major limb trauma. Ann Surg 2009; 249:1. Walters TJ, Wenke JC, Kauvar DS, et al. Effectiveness of self-applied tourniquets in human volunteers. Prehosp Emerg Care 2005; 9:416. Inaba K, Siboni S, Resnick S, et al. Tourniquet use for civilian extremity trauma. J Trauma Acute Care Surg 2015; 79:232. Schroll R, Smith A, McSwain NE Jr, et al. A multi-institutional analysis of prehospital tourniquet use. J Trauma Acute Care Surg 2015; 79:10. Leonard J, Zietlow J, Morris D, et al. A multi-institutional study of hemostatic gauze and tourniquets in rural civilian trauma. J Trauma Acute Care Surg 2016; 81:441. US Department of Health and Human Services Centers for Disease Control and Prevention. Tetanus. www.cdc.gov/vaccines/pubs/pinkbook/downloads/tetanus.pdf (Accessed on May 06, 2011). Cavadas PC, LandÃn L, IbÃjñez J, et al. Infrapopliteal lower extremity replantation. Plast Reconstr Surg 2009; 124:532. Sapega AA, Heppenstall RB, Sokolow DP, et al. The bioenergetics of preservation of limbs before replantation. The rationale for intermediate hypothermia. J Bone Joint Surg Am 1988; 70:1500. Cancio LC, Jimenez-Reyna JF, Barillo DJ, et al. One hundred ninety-five cases of high-voltage electric injury. J Burn Care Rehabil 2005; 26:331. Bosse MJ, McCarthy ML, Jones AL, et al. The insensate foot following severe lower extremity trauma: an indication for amputation? J Bone Joint Surg Am 2005; 87:2601. Burkhardt GE, Cox M, Clouse WD, et al. Outcomes of selective tibial artery repair following combat-related extremity injury. J Vasc Surg 2010; 52:91. Lynch K, Johansen K. Can Doppler pressure measurement replace "exclusion" arteriography in the diagnosis of occult extremity arterial trauma? Ann Surg 1991; 214:737. Mills WJ, Barei DP, McNair P. The value of the ankle-brachial index for diagnosing arterial injury after knee dislocation: a prospective study. J Trauma 2004; 56:1261. Wallin D, Yaghoubian A, Rosing D, et al. Computed tomographic angiography as the primary diagnostic modality in penetrating lower extremity vascular injuries: a level I trauma experience. Ann Vasc Surg 2011; 25:620. Soto JA, Múnera F, Cardoso N, et al. Diagnostic performance of helical CT angiography in trauma to large arteries of the extremities. J Comput Assist Tomogr 1999; 23:188. Soto JA, Múnera F, Morales C, et al. Focal arterial injuries of the proximal extremities: helical CT arteriography as the initial method of diagnosis. Radiology 2001; 218:188. Jens S, Kerstens MK, Legemate DA, et al. Diagnostic performance of computed tomography angiography in peripheral arterial injury due to trauma: a systematic review and meta-analysis. Eur J Vasc Endovasc Surg 2013; 46:329. Melvin JS, Dombroski DG, Torbert JT, et al. Open tibial shaft fractures: I. Evaluation and initial wound management. J Am Acad Orthop Surg 2010; 18:10. Seamon MJ, Smoger D, Torres DM, et al. A prospective validation of a current practice: the detection of extremity vascular injury with CT angiography. J Trauma 2009; 67:238. White PW, Gillespie DL, Feurstein I, et al. Sixty-four slice multidetector computed tomographic angiography in the evaluation of vascular trauma. J Trauma 2010; 68:96. Propper BW, Alley JB, Gifford SM, et al. Endovascular treatment of a blunt aortic injury in Iraq: extension of innovative endovascular capabilities to the modern battlefield. Ann Vasc Surg 2009; 23:687.e19. Yan H, Gao W, Li Z, et al. The management of degloving injury of lower

extremities: technical refinement and classification. J Trauma Acute Care Surg 2013; 74:604. Harboe K, SÃ, reide K. Tibial fracture with degloving injury of the foot. Br J Surg 2012; 99 Suppl 1:87. Nickerson TP, Zielinski MD, Jenkins DH, Schiller HJ. The Mayo Clinic experience with Morel-Lavallée lesions: establishment of a practice management guideline. J Trauma Acute Care Surg 2014; 76:493. Konda SR, Davidovitch RI, Egol KA. Computed tomography scan to detect traumatic arthrotomies and identify periarticular wounds not requiring surgical intervention: an improvement over the saline load test. J Orthop Trauma 2013; 27:498. Konda SR, Howard D, Davidovitch RI, Egol KA. The role of computed tomography in the assessment of open periarticular fractures associated with deep knee wounds. J Orthop Trauma 2013; 27:509. Nord RM, Quach T, Walsh M, et al. Detection of traumatic arthrotomy of the knee using the saline solution load test. J Bone Joint Surg Am 2009; 91:66. Keese GR, Boody AR, Wongworawat MD, Jobe CM. The accuracy of the saline load test in the diagnosis of traumatic knee arthrotomies. J Orthop Trauma 2007; 21:442. Konda SR, Howard D, Davidovitch RI, Egol KA. The saline load test of the knee redefined: a test to detect traumatic arthrotomies and rule out periarticular wounds not requiring surgical intervention. J Orthop Trauma 2013; 27:491. Owens BD, Kragh JF Jr, Macaitis J, et al. Characterization of extremity wounds in Operation Iragi Freedom and Operation Enduring Freedom. J Orthop Trauma 2007; 21:254. Gustilo RB, Mendoza RM, Williams DN. Problems in the management of type III (severe) open fractures: a new classification of type III open fractures. J Trauma 1984; 24:742. Gustilo RB, Anderson JT. Prevention of infection in the treatment of one thousand and twenty-five open fractures of long bones: retrospective and prospective analyses. J Bone Joint Surg Am 1976; 58:453. Okike K, Bhattacharyya T. Trends in the management of open fractures. A critical analysis. J Bone Joint Surg Am 2006; 88:2739. Brumback RJ, Jones AL. Interobserver agreement in the classification of open fractures of the tibia. The results of a survey of two hundred and forty-five orthopaedic surgeons. J Bone Joint Surg Am 1994; 76:1162. Orthopaedic Trauma Association: Open Fracture Study Group. A new classification scheme for open fractures. J Orthop Trauma 2010; 24:457. Patterson BM, Agel J, Swiontkowski MF, et al. Knee dislocations with vascular injury: outcomes in the Lower Extremity Assessment Project (LEAP) Study. J Trauma 2007; 63:855. Dua A, Desai SS, Shah JO, et al. Outcome predictors of limb salvage in traumatic popliteal artery injury. Ann Vasc Surg 2014; 28:108. American College of Surgeons Committee on Trauma ad hoc Committee on Outcomes. Management of Complex Extremity Trauma. www.facs.org/trauma/publications/mancompexttrauma.pdf (Accessed on May 06, 2011). Arrillaga A, Bynoe R, Frykberg ER, Nagy K. EAST Practice Management Guidelines for Penetrating Trauma to the Lower Extremity. http://www.east.org/Portal/Default.aspx?tabid=57 (Accessed on May 06, 2011). Scalea JR, Crawford R, Scurci S, et al. Below-the-knee arterial injury: the type of vessel may be more important than the number of vessels injured. J Trauma Acute Care Surg 2014; 77:920. Branco BC, Linnebur M, Boutrous ML, et al. The predictive value of multidetector CTA on outcomes in patients with below-the-knee vascular injury. Injury 2015; 46:1520. Bosse MJ, MacKenzie EJ, Kellam JF, et al. A prospective evaluation of the clinical utility of the lower-extremity injury-severity scores. J Bone Joint Surg Am 2001; 83-A:3. Alarhayem AQ, Cohn SM, Cantu-Nunez O, et al. Impact of time to repair on outcomes in patients with lower extremity arterial injuries. J Vasc Surg 2019; 69:1519. Harris AM, Althausen PL, Kellam J, et al. Complications following limb-threatening lower extremity trauma. J Orthop Trauma 2009;

23:1. <u>Rozycki GS, Tremblay LN, Feliciano DV, McClelland WB. Blunt vascular trauma in the</u> extremity: diagnosis, management, and outcome. J Trauma 2003; 55:814.

Gibbons RJ, Balady GJ, Bricker JT, et al. ACC/AHA 2002 guideline update for exercise testing: summary article. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1997 Exercise Testing Guidelines). J Am Coll Cardiol 2002; 40:1531. Hendel RC, Berman DS, Di Carli MF, et al. ACCF/ASNC/ACR/AHA/ASE/SCCT/SCMR/SNM 2009 Appropriate Use Criteria for Cardiac Radionuclide Imaging: A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, the American Society of Nuclear Cardiology, the American College of Radiology, the American Heart Association, the American Society of Echocardiography, the Society of Cardiovascular Computed Tomography, the Society for Cardiovascular Magnetic Resonance, and the Society of Nuclear Medicine. J Am Coll Cardiol 2009; 53:2201. American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Society of Echocardiography, American Heart Association, et al. ACCF/ASE/AHA/ASNC/HFSA/HRS/SCAI/SCCM/SCCT/SCMR 2011 Appropriate Use Criteria for Echocardiography. A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Society of Echocardiography, American Heart Association, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance American College of Chest Physicians. J Am Soc Echocardiogr 2011; 24:229. Pellikka PA, Nagueh SF, Elhendy AA, et al. American Society of Echocardiography recommendations for performance, interpretation, and application of stress echocardiography. J Am Soc Echocardiogr 2007; 20:1021. Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. J Am Coll Cardiol 2012; 60:e44. American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, American Society of Echocardiography, American Society of Nuclear Cardiology, et al. 2009 ACCF/AHA focused update on perioperative beta blockade. J Am Coll Cardiol 2009; 54:2102. Wolk MJ, Bailey SR, Doherty JU, et al. ACCF/AHA/ASE/ASNC/HFSA/HRS/SCAI/SCCT/SCMR/STS 2013 multimodality appropriate use criteria for the detection and risk assessment of stable ischemic heart disease: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. J Am Coll Cardiol 2014; 63:380. Miller TD. Exercise treadmill test: estimating cardiovascular prognosis. Cleve Clin J Med 2008; 75:424. Kligfield P, Lauer MS. Exercise electrocardiogram testing: beyond the ST segment. Circulation 2006; 114:2070. Mark DB, Hlatky MA, Harrell FE Jr,

et al. Exercise treadmill score for predicting prognosis in coronary artery disease. Ann Intern Med 1987; 106:793. Mark DB, Shaw L, Harrell FE Jr, et al. Prognostic value of a treadmill exercise score in outpatients with suspected coronary artery disease. N Engl J Med 1991; 325:849. Skopicki HA, Abraham SA, Picard MH, et al. Effects of dobutamine at maximally tolerated dose on myocardial blood flow in humans with ischemic heart disease. Circulation 1997; 96:3346. Tadamura E, Iida H, Matsumoto K, et al. Comparison of myocardial blood flow during dobutamine-atropine infusion with that after dipyridamole administration in normal men. J Am Coll Cardiol 2001; 37:130. Bateman TM, Heller GV, McGhie AI, et al. Diagnostic accuracy of rest/stress ECG-gated Rb-82 myocardial perfusion PET: comparison with ECG-gated Tc-99m sestamibi SPECT. J Nucl Cardiol 2006; 13:24. Arruda-Olson AM, Juracan EM, Mahoney DW, et al. Prognostic value of exercise echocardiography in 5,798 patients: is there a gender difference? J Am Coll Cardiol 2002; 39:625. Heitner JF, Kim RJ, Kim HW, et al. Prognostic Value of Vasodilator Stress Cardiac Magnetic Resonance Imaging: A Multicenter Study With 48â€[−]000 Patient-Years of Follow-up. JAMA Cardiol 2019; 4:256. Fleischmann KE, Hunink MG, Kuntz KM, Douglas PS. Exercise echocardiography or exercise SPECT imaging? A meta-analysis of diagnostic test performance. JAMA 1998; 280:913. Garber AM, Solomon NA. Cost-effectiveness of alternative test strategies for the diagnosis of coronary artery disease. Ann Intern Med 1999; 130:719. Gianrossi R, Detrano R, Mulvihill D, et al. Exercise-induced ST depression in the diagnosis of coronary artery disease. A meta-analysis. Circulation 1989; 80:87. Chareonthaitawee P, Gersh BJ, Araoz PA, Gibbons RJ. Revascularization in severe left ventricular dysfunction: the role of viability testing. J Am Coll Cardiol 2005; 46:567. O'Rourke RA, Brundage BH, Froelicher VF, et al. American College of Cardiology/American Heart Association Expert Consensus document on electron-beam computed tomography for the diagnosis and prognosis of coronary artery disease. Circulation 2000; 102:126. Detrano R, Hsiai T, Wang S, et al. Prognostic value of coronary calcification and angiographic stenoses in patients undergoing coronary angiography. J Am Coll Cardiol 1996; 27:285. Keelan PC, Bielak LF, Ashai K, et al. Long-term prognostic value of coronary calcification detected by electron-beam computed tomography in patients undergoing coronary angiography. Circulation 2001; 104:412. Douglas PS, Hoffman U, Patel MR, et al.. Outcomes of anatomical versus functional testing for coronary artery disease. N Engl J Med 2015; 372:1291. Gongora CA, Bavishi C, Uretsky S, Argulian E. Acute chest pain evaluation using coronary computed tomography angiography compared with standard of care: a meta-analysis of randomised clinical trials. Heart 2018; 104:215. Truong QA, Schulman-Marcus J, Zakroysky P, et al. Coronary CT Angiography Versus Standard Emergency Department Evaluation for Acute Chest Pain and Diabetic Patients: Is There Benefit With Early Coronary CT Angiography? Results of the Randomized Comparative Effectiveness ROMICAT II Trial. J Am Heart Assoc 2016; 5:e003137. Hamilton-Craig C, Fifoot A, Hansen M, et al. Diagnostic performance and cost of CT angiography versus stress ECG--a randomized prospective study of suspected acute coronary syndrome chest pain in the emergency department (CT-COMPARE). Int J Cardiol 2014; 177:867. SCOT-HEART investigators. CT coronary angiography in patients with suspected angina due to coronary heart disease (SCOT-HEART): an open-label, parallel-group, multicentre trial. Lancet 2015; 385:2383. SCOT-HEART Investigators, Newby DE, Adamson PD, et al. Coronary CT Angiography and 5-Year Risk of Myocardial Infarction. N Engl J Med 2018; 379:924. Lubbers M, Dedic A, Coenen A, et al. Calcium imaging and selective computed tomography angiography in

<u>comparison to functional testing for suspected coronary artery disease: the multicentre,</u> <u>randomized CRESCENT trial. Eur Heart J 2016; 37:1232.</u> <u>Lubbers M, Coenen A, Kofflard M, et al.</u> <u>Comprehensive Cardiac CT With Myocardial Perfusion Imaging Versus Functional Testing in</u> <u>Suspected Coronary Artery Disease: The Multicenter, Randomized CRESCENT-II Trial. JACC</u> <u>Cardiovasc Imaging 2018; 11:1625.</u>